

(8 pages)

Reg. No. :

Code No. : 10427 E Sub. Code : CSMA 41

B.Sc. (CBCS) DEGREE EXAMINATION, APRIL 2023

Fourth Semester

Mathematics — Skill Based Subject

TRIGONOMETRY, LAPLACE TRANSFORMS AND
FOURIER SERIES

(For those who joined in July 2021 onwards)

Time : Three hours

Maximum : 75 marks

PART A — (10 × 1 = 10 marks)

Answer ALL questions.

Choose the correct answer :

1. 1 radian = _____ degree.
(a) 52.79 (b) 57.29
(c) 59.27 (d) 59.29
2. If $\tan \theta = \frac{1}{15}$ then $\theta \simeq$ _____
(a) $3^\circ 47'$ (b) $3^\circ 41'$
(c) $3^\circ 49'$ (d) $3^\circ 94'$

3. The value of $\tanh^{-1} x$ is _____

- (a) $\frac{1}{2} \log \left(\frac{1+x}{1-x} \right)$ (b) $\frac{1}{2} \log \left(\frac{1-x}{1+x} \right)$
(c) $\frac{1}{2} \log \left(\frac{x-1}{x+1} \right)$ (d) $\frac{1}{2} \log \left(\frac{x+1}{x-1} \right)$

4. The value of $i^i =$ _____

- (a) $e^{\frac{(4n-1)\pi}{2}}$ (b) $e^{\frac{-(4n+1)\pi}{2}}$
(c) $e^{\frac{-(4n-1)\pi}{2}}$ (d) $e^{(4n-1)\pi}$

5. $L[f'(x)] =$ _____

- (a) $f(0) - SL[f(x)]$
(b) $SL[f(x)] - f(0)$
(c) $f'(0) - SL[f(x)]$
(d) $S^2 L[f(x)] - Sf(0) - f'(0)$

6. $L^{-1}[F(s+a)] =$ _____

- (a) $e^{-ax} L^{-1}[f(s)]$ (b) $e^{ax} L^{-1}[f(s)]$
(c) $e^{ax} L[f(s)]$ (d) $\frac{1}{a} f\left(\frac{s}{a}\right)$



7. $L[xy''] = \text{—————}$

(a) $\frac{d}{dS}(S^2 L(y) - Sy'(0) - y'(0))$

(b) $\frac{d}{dS}(S^2 L(y) - Sy'(0) - y(0))$

(c) $-\frac{d}{dS}(S^2 L(y) - Sy'(0) - y'(0))$

(d) $-\frac{d}{dS}(S^2 L(y) - Sy'(0) - y(0))$

8. $L(xy') = \text{—————}$

(a) $-\frac{d}{dS}[SL(y) - y(0)]$

(b) $\frac{d}{dS}[SL(y) - y(0)]$

(c) $\frac{d}{dS}[SL(y) - y'(0)]$

(d) $-\frac{d}{dS}[SL(y) - y'(0)]$

9. The Fourier co-efficient a_0 for $f(x) = x^2$ in $(-\pi, \pi)$ is ———

(a) 0 (b) $\frac{2\pi^3}{3}$

(c) $\frac{2\pi^2}{3}$ (d) $\frac{\pi^3}{3}$

10. For any integer n , the value of $\cos n\pi$ is ———

(a) 1 (b) 0

(c) -1 (d) $(-1)^n$

PART B — (5 × 5 = 25 marks)

Answer ALL questions choosing either (a) or (b).

11. (a) Prove that

$$2^5 \cos^6 \theta = \cos 6\theta + 6 \cos 4\theta + 15 \cos 2\theta + 10.$$

Or

(b) Evaluate $\lim_{x \rightarrow \frac{\pi}{2}} \left[\frac{\sin x + \cos 2x}{\cos^2 x} \right]$.



12. (a) If $\cos(x + iy) = r(\cos \alpha + i \sin \alpha)$. Then prove that $y = \frac{1}{2} \log \left[\frac{\sin(x - \alpha)}{\sin(x + \alpha)} \right]$.

Or

- (b) Sum to infinity the series $1 + \frac{c^2 \cos 2\theta}{2!} + \frac{c^4 \cos 4\theta}{4!} + \dots + \infty$.

13. (a) Find $L \left[\frac{1 - \cos x}{x} \right]$.

Or

- (b) Find $L^{-1} \left[\log \left(\frac{s+a}{s+b} \right) \right]$.

14. (a) Using Laplace transform, solve $y' + 3y = e^{-2x}$ given $y(0) = 4$.

Or

- (b) Solve $(D^2 + 5D + 6)y = e^{-x}$ given that $y(0) = 0$ and $y'(0) = 0$, using Laplace transform.

15. (a) If $f(x) = \begin{cases} -\pi/4 & \text{if } -\pi < x < 0 \\ \pi/4 & \text{if } 0 < x < \pi \end{cases}$, then find the Fourier series of $f(x)$.

Or

- (b) Find the Fourier constant b_1 , for the function $x \sin x$ in the half range $0 < x < \pi$.

PART C — (5 × 8 = 40 marks)

Answer ALL questions choosing either (a) or (b).

16. (a) Prove that $\frac{\cos 9\theta}{\cos \theta} = 256 \cos^8 \theta - 576 \cos^6 \theta + 432 \cos^4 \theta - 120 \cos^2 \theta + 9$.

Or

- (b) Prove that

$$\cos^5 \theta \sin^4 \theta = \left(\frac{1}{2} \right)^8 [\cos 9\theta + \cos 7\theta - 4 \cos 5\theta - 4 \cos 3\theta + 6 \cos \theta].$$

17. (a) If $\tan(\theta + i\phi) = \cos \alpha + i \sin \alpha$, prove that

$$(i) \quad \theta = \frac{1}{2} n\pi + \frac{\pi}{4}$$

$$(ii) \quad \phi = \frac{1}{2} \log \tan \left(\frac{\pi}{4} + \frac{\alpha}{2} \right).$$

Or



- (b) Sum the series to infinity

$$x \sin \theta + \frac{x^3}{3} \sin 3\theta + \frac{x^5}{5} \sin 5\theta + \dots + \infty.$$

18. (a) Find

(i) $L[t^2 + \cos 2t \cos t + \sin^2 2t]$

(ii) $L[xe^{-x} \cos x].$

Or

- (b) Find

(i) $L^{-1}\left[\frac{s}{(s+2)^2}\right]$

(ii) $L^1\left[\frac{s^2}{(s-1)^2}\right].$

19. (a) Solve $y'' - 4y' + 4y = x$ given that $y(0) = 0$ and $y'(0) = 1$ using Laplace transform.

Or

- (b) Using Laplace transform, solve the equation $xy'' - (2x+1)y' + (x+1)y = 0$ given that $y(0) = 0$.

Page 7 Code No. : 10427 E

20. (a) Find the Fourier series of $f(x) = |\sin x|$ in $(-\pi, \pi)$ of periodicity 2π .

Or

- (b) Prove that the function $f(x) = x$ can be expressed in a series of cosine in $0 \leq x \leq \pi$ as

$$x = \frac{\pi}{2} - \frac{\pi}{4} \left[\frac{\cos x}{1^2} + \frac{\cos 3x}{3^2} + \frac{\cos 5x}{5^2} + \dots \right]. \text{ Hence}$$

$$\text{deduce that } \frac{1}{1^2} + \frac{1}{3^2} + \dots = \frac{\pi^2}{8}.$$

Page 8 Code No. : 10427 E

