(6 pages)	Reg. No. :		3.	Let f be measur	able. inf	$\{\alpha \mid f \leq \alpha \ a.e\}$	is called
Code No. : 5	761 Sub. Code :	WMAM 22		(a) essential supr	emum		
M.Sc. (CBCS) DEGREE EXAMINATION, APRIL 2024.				(b) supermum (c) essential infimum			
Second Semester				(d) balanced infimum			
Mathematics — Core REAL ANALYSIS — II			4.	If f is a Riemann integral over $[a, b]$ then $\int_{a}^{b} f$			
(For those who joined in July 2023 onwards)				$\qquad \qquad \int_a^b f .$			
Time : Three hou	rs Maximun	n: 75 marks		(a) <	(b)	=	
PAF	$TA - (15 \times 1 = 15 \text{ marks})$			(c) >	(d)	≠	
Answer ALL questions. Choose the correct answer:			5.	If f is integrable th	nen $\left \int f dx \right $		$\int f dx$.
1. The measu	re of an Interval I, denoted	d by $m(I) =$		(a) <	(b)	<	
(a) Initial	 value (b) Terminal	value		(c) ≥	(d)	>	
(a) not me	2. Continuous functions are (a) not measurable		6.	A non negative infinite valued function taking only finite number of values is calledfunction.			
(b) absolutely measurable (c) measurable				(a) finite	(b)	measurable	9
	v measurable			(c) constant	(b)	simple	

Page 2

Code No. : 5761

- The Inner Product of f, g is $\langle f, g \rangle =$ _
 - (a) $\int \overline{f(x)}g(x)dx$ (b) $\int g(x)dx$
- - (c) $\int f(x)\overline{g(x)}dx$ (d) $\int f(x)dx$
- $\lim_{n\to\infty}\int_{0}^{2\pi}f(x)\sin nxdx = --$
 - (a) 0
- (b) 2
- (c) 1

- (d) π
- A function ϕ_n is orthonormal if $\|\phi_n\|$ 9.
 - (a) 0

(b) -1

(c) 2

- (d) 1
- 10. A function $f: \mathbb{R}^n \to \mathbb{R}^m$ is linear if f(ax + by) =
 - (a) af(x)+bf(y) (b) af(x)-bf(y)

 - (c) bf(x) af(y) (d) bf(x) + af(y)
- 11. A function f is differentiable at c if there exists $T_c: \mathbb{R}^n \to \mathbb{R}^m$ such that $f(c+v) = \underline{}$

 - (a) $T_c |v| + ||v||$ (b) $f(c) + T_c |v| + ||v|| E_c(v)$
 - (c) $T_c|v| ||v|| E_c(v)$ (d) $||v|| E_c(v)$

Code No.: 5761 Page 3

- 12. If f is a linear function then f(c+v)=
 - (a) f(c) f(v) (b) f(v) f(c)

- (c) f(v)
- (d) f(c)+f(v)
- 13. The set [0, 1] is _____
 - (a) countable
- (b) non bounded
- (c) bounded
- (d) not countable
- 14. A function f has local extremum at c if,
 - (a) f'(c) > 0
- (b) f'(c) = 0
- (c) f'(c) < 0
- (d) $f'(c) \neq 0$
- 15. A function $f: s \to T$ from (S, d_S) to (T, d_T) is an open mapping if for A in S, f(A) is _____ in
 - (a) open

- (b) closed
- (c) imbedded
- (d) bounded

PART B —
$$(5 \times 4 = 20 \text{ marks})$$

Answer ALL questions by choosing either (a) or (b).

(a) If $m^*(A)$ is finite show that $m^*(\phi) = 0$.

Or

(b) Show that a countable set has outer measure zero.

Page 4

Code No.: 5761

[P.T.O.]

17. (a) Prove Lebesgue dominated convergence theorem.

Or

- (b) State and prove Simple Approximation Lemma.
- 18. (a) Prove chain rule among functions.

Or

- (b) State Riemann Localization Theorem.
- 19. (a) Let $f: S \to \mathbb{R}^n$. Let C be an interior point of S. If f is differentiable at C then show that f is continuous at C.

Or

- (b) Derive Parseval's Formula.
- 20. (a) Prove Bounded Convergence Theorem.

Or

(b) Prove Egoroff's theorem.

PART C —
$$(5 \times 8 = 40 \text{ marks})$$

Answer ALL questions by choosing either (a) or (b).

21. (a) Prove that not every measurable set is a Borel Set.

Or

(b) Prove the outer measure of intervals is its length.

Page 5 Code No.: 5761

22. (a) Prove monotone convergence theorem on measurable functions.

Or

- (b) State and prove Fatou's Lemma.
- 23. (a) Prove Riemann-Lebesgue Lemma.

Or

- (b) State and prove Riesz-Fischer Theorem.
- 24. (a) Prove Taylor's Formula.

Or

- (b) Prove mean value theorem for differentiable functions.
- 25. (a) State and prove Implicit Function Theorem.

Or

(b) State and prove Vitali's Covering Lemma.

Page 6 Code No.: 5761